



Australian Government



After your COVID-19 vaccination

About the vaccine

Today you have received the Comirnaty (Pfizer Australia Pty Ltd) vaccine.

This vaccine can prevent people from becoming ill from COVID-19. Comirnaty does not contain any live virus, and it cannot give you COVID-19. It contains the genetic code for an important part of the SARS-CoV-2 virus called the spike protein. After getting the vaccine, your body makes copies of the spike protein, and your immune system will learn to recognise and fight against the SARS-CoV-2 virus.

What to expect after vaccination

As with any vaccine, you may have some side effects after receiving a COVID-19 vaccine.

Common side effects after **Comirnaty** include:

- pain or swelling at the injection site
- tiredness
- headache
- muscle pain
- chills
- fever
- joint pain.

Less common side effects after **Comirnaty** include:

- redness at the injection site
- nausea
- enlarged lymph nodes
- feeling unwell
- pain in limb
- insomnia
- itching at the injection site.



These side effects are usually mild and usually go away within one or two days. If you experience pain at the injection site or fever, headaches or body aches after vaccination, you can take paracetamol or ibuprofen. These help reduce the above symptoms (you do not need to take paracetamol or ibuprofen before vaccination). If there is swelling at the injection site, you can use a cold compress.

Rare side effects that have been reported after **Comirnaty** are:

- Severe allergic reaction (anaphylaxis)
- Temporary one-sided facial drooping (also called Bell's palsy).

You should seek medical attention after vaccination if:

- You think you are having an allergic reaction. Call 000 if you experience severe symptoms, such as difficulty breathing, wheezing, a fast heartbeat or collapsing.
- You are worried about a potential side effect or have new or unexpected symptoms.
- You have an expected side effect of the vaccine which has not gone away after a few days.

For symptoms which are not urgent, you can see your regular healthcare provider (usually your GP).

Vaccine safety monitoring and reporting side effects

Suspected side effects can be reported to your vaccination provider or other healthcare professional. They will then make a formal report on your behalf to your state or territory health department or directly to the Therapeutic Goods Administration (TGA).

If you would prefer to report it yourself, please visit the [reporting suspected side effects associated with a COVID-19 vaccine webpage](#) on the TGA website and follow the directions on the page.

COVID-19 testing after vaccination

Some side effects from COVID-19 vaccination might be similar to symptoms of COVID-19 (e.g. fever), though Comirnaty does not contain any live virus and cannot cause COVID-19. You may not need to get a COVID-19 test or isolate if you develop general symptoms like fever, headache or tiredness in the first two days after vaccination and if you are sure that you don't have any respiratory symptoms (e.g. runny nose, cough, sore throat, loss of smell or loss of taste). However, you should check the current guidelines in your state/territory for the most up-to-date information. This advice may change in the event of a COVID-19 outbreak in your local area. You may still need to get a COVID-19 test if you meet other criteria, e.g. you are a close contact of a known COVID-19 case. If in doubt, seek medical assessment.

Remember your second appointment

It is important that you receive two doses of Comirnaty (Pfizer), at least three weeks apart. Maximum protection against COVID-19 will not occur until at least a week after your second dose.

How is the information you provide at your appointment used

For information on how your personal details are collected, stored and used visit <https://www.health.gov.au/covid19-vaccines>